



MARILYN'S MAKE UP MAGIC

@ New Image Plastic Surgery
7425 e. Shea Blvd., Ste. 105
Scottsdale, AZ 85260
480-947-8498

PRE-TREATMENT CARE

Health conditions and medications that slow down healing, cause swelling, bleeding and loss of tattoo color are:

Alcohol Consumption	High Blood Pressure	Aspirin
Allergies	Sun Damage	Ibuprofen
Diabetes	Scar Tissue	Anti-inflammatories
Cold Sores	Skin Disorders	Decongestants
Frequent Infections	Vitamin A & E (mega-doses)	Dermabrasion
Pre-cancer	Garlic	Omega 3

Discontinue mega-doses of Vitamin A, Vitamin E, Omega 3, Garlic and all non-prescription medications (except Tylenol), five days prior to procedure. Reduce salt intake and discontinue ALL ALCOHOLIC BEVERAGES at least 24 hours before procedure. **PLEASE DO NOT WEAR ANY PERFUME.**

PLEASE CONSULT YOUR DOCTOR before discontinuing any **prescribed** medication(s) or before taking narcotic, prescription pain relievers, especially if you have any of the following: PREGNANCY, MITRO VALVE PROLAPSE, HEART MURMUR, RHEUMATIC FEVER, JOINT REPLACEMENT (S), PACEMAKER, recent OPEN HEART SURGERY, or remaining PLATES OR PINS IN BONES. Any of the preceding may require pre-medication (i.e., antibiotics) or other precautions recommended by your physician.

EYEBROW PROCEDURE: Brows can appear dark for 3-7 days. Minimal redness and irritation can be expected during this time.

EYELINER PROCEDURE: Consider contacting your doctor regarding a prescribed **NARCOTIC pain reliever** to take during each eyeliner appointment. **This is to relax the eye muscles to control eye muscle spasms – NOT PAIN!** For this procedure, most doctors prescribe **Codeine, Percocet, Roxicet**, etc. Bring the pain reliever to your appointment. You may want to consider a prescription for an ophthalmic-steroid eye ointment, for use at home, which is beneficial to reduce swelling if used immediately following this procedure. (Non-ophthalmic Bacitracin is provided during the procedure.) You should have **ARTIFICIAL TEARS or OPTHALMIC SALINE SOLUTION and ophthalmic ointment available for home care treatment.**

LIP CORRECTION PROCEDURE: If you have **ever, ever, ever** had a cold sore outbreak, you still have the virus in your system! Lip tissue damage is an open invitation for this viral infection resulting in swelling, pain and causing some color loss in your new lip tattoo! Your Physician will discuss an **oral** prescription, (i.e., **Valtrex & Denavir**) to prevent an outbreak. Most doctors recommend 1000 mg .of an oral, anti-viral medication for 5 days, beginning the day before the lip procedure. This routine should be repeated on all follow-up procedures. The topical prescription ointment called Denavir is very helpful in preventing/controlling cold sore outbreaks.

Lips will be anesthetized by prescription medical-grade topical numbing compound. **NO INJECTIONS! NO PAIN!!**

SWELLING is unpredictable and varies with each individual. However, swelling is reduced substantially for most individuals in 24-48 hours. The injections may aggravate soft tissue in some individuals, thereby increasing swelling; others will swell with or without injection. Color is quite dark during the swollen period, but mellows as swelling subsides. It helps to keep head elevated above heart for the first 24 hours.

Lips will be more receptive to color and have less tissue damage if softened with daily massage of **"VITAMIN E" OIL** (for two weeks or more) before each lip procedure. Do not use Vitamin E after lip tattoo until it is completely healed.

For individuals with frequent viral outbreaks, please contact your doctor regarding a preventive program.

NOTE → → → → NO PERFUME PLEASE! MARILYN IS ALLERGIC ☹

THERE WILL BE A CHARGE FOR EACH PROCEDURE FOR ANY FOLLOW UP APPOINTMENTS.

CASH OR CHECK - NO CREDIT CARDS. ALL PRICES ARE SUBJECT TO CHANGE.

- No guarantees or assurances have been given to me regarding this procedure.
- I understand that perfection is not a realistic expectation.
- I understand that numerous treatments may be necessary to obtain the desired result and that there is a charge required for any additional treatments.

Non-Ophthalmic Bacitracin is applied during and after procedure. Please let the technician know if you are sensitive to Neosporin, Polysporin, Bacitracin or other petroleum based products. Secondary swelling or inflammation is possible. If sensitivity, excessive irritation, swelling or redness develops, discontinue usage and use warm compresses every 1-2 hours until swelling and inflammation is gone. It is also very important that you let our office know of any allergic reactions to any topicals used.